



Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

# Introduction to Network Analysis

Structure of networks: cohesion

Vladimir Batagelj

IMFM Ljubljana, IAM UP Koper and NRU HSE Moscow

**Master's programme**

**Applied Statistics with Social Network Analysis**

International Laboratory for Applied Network Research

NRU HSE, Moscow 2018

Rnet,  
cohesion


V. Batagelj

Islands

Cores

Generalized  
cores

- 1 Islands
- 2 Cores
- 3 Generalized cores




L. Krempf, MPI.

Vladimir Batagelj: [vladimir.batagelj@fmf.uni-lj.si](mailto:vladimir.batagelj@fmf.uni-lj.si)

Current version of slides (December 11, 2018 at 12:18): [slides PDF](#)

If we represent a given or computed value of nodes / links as a height of nodes / links and we immerse the network into a water up to selected level we get *islands*. Varying the level we get different islands.



We developed very efficient algorithms to determine the islands hierarchy and to list all the islands of selected sizes.  
See [details](#).



## ... Islands

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

Islands are very general and efficient approach to determine the 'important' subnetworks in a given network.

We have to express the goals of our analysis with a related property of the nodes or weight of the links. Using this property we determine the islands of an appropriate size (in the interval  $k$  to  $K$ ).

In large networks we can get many islands which we have to inspect individually and interpret their content.

An important property of the islands is that they identify locally important subnetworks on different levels. Therefore they detect also emerging groups.

The set of nodes  $\mathcal{C} \subseteq \mathcal{V}$  is a **local node peak**, if it is a regular node island and all of its nodes have the same value. Node island with a single local node peak is called a **simple node island**. In similar way we define simple link island.



## ... Islands

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

A set of nodes  $C \subseteq \mathcal{V}$  is a *regular node island* in network  $\mathcal{N} = (\mathcal{V}, \mathcal{L}, p)$ ,  $p : \mathcal{V} \rightarrow \mathbb{R}$  iff it induces a connected subgraph and the nodes from the island are 'higher' than the neighboring nodes

$$\max_{u \in N(C)} p(u) < \min_{v \in C} p(v)$$

A set of nodes  $C \subseteq \mathcal{V}$  is a *regular link island* in network  $\mathcal{N} = (\mathcal{V}, \mathcal{L}, w)$ ,  $w : \mathcal{L} \rightarrow \mathbb{R}$  iff it induces a connected subgraph and the links inside the island are 'stronger related' among them than with the neighboring nodes – in  $\mathcal{N}$  there exists a spanning tree  $\mathcal{T}$  over  $C$  such that

$$\max_{(u,v) \in \mathcal{L}, u \notin C, v \in C} w(u, v) < \min_{(u,v) \in \mathcal{T}} w(u, v)$$

Network/Create Partition/Islands/Line Weights  
Operations/Network+Vector/Islands/Vertex  
Property





# Some properties of node islands

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

- The sets of nodes of connected components of node-cut at selected level  $t$  are regular node islands.
- The set  $\mathcal{H}_p(\mathcal{N})$  of all regular node islands of network  $\mathcal{N}$  is a complete hierarchy:
  - two islands are disjoint or one of them is a subset of the other
  - each node belongs to at least one island
- Node islands are invariant for the strictly increasing transformations of the property  $p$ .
- Two linked nodes cannot belong to two disjoint regular node islands.



# Simple node islands

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

- The set of nodes  $\mathcal{C} \subseteq \mathcal{V}$  is a **local node peak**, if it is a regular node island and all of its nodes have the same value.
- Node island with a single local node peak is called a **simple node island**.
- The types of node islands:
  - FLAT – all nodes have the same value
  - SINGLE – island has a single local node peak
  - MULTI – island has more than one local node peaks
- Only the islands of type FLAT or SINGLE are simple islands.



# Some properties of link islands

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

- The sets of nodes of connected components of link-cut at selected level  $t$  are regular link islands.
- The set  $\mathcal{H}_w(\mathcal{N})$  of all nondegenerated regular link islands of network  $\mathcal{N}$  is hierarchy (not necessarily complete):
  - two islands are disjoint or one of them is a subset of the other
- Link islands are invariant for the strictly increasing transformations of the weight  $w$ .
- Two linked nodes may belong to two disjoint regular link islands.





# Simple link islands

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

- The set of nodes  $\mathcal{C} \subseteq \mathcal{V}$  is a **local link peak**, if it is a regular link island and there exists a spanning tree of the corresponding induced network, in which all links have the same value as the link with the largest value.
- Link island with a single local link peak is called a **simple link island**.
- The types of link islands:
  - FLAT – there exists a spanning tree, in which all links have the same value as the link with the largest value.
  - SINGLE – island has a single local link peak.
  - MULTI – island has more than one local link peaks.
- Only the islands of type FLAT or SINGLE are simple islands.



# US patents

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

US patents network (Nber, US Patents) has 3774768 nodes and 16522438 arcs (1 loop). Without the loop it is acyclic. The weight of an arc is the proportion of paths through the arc from some initial node to some terminal node. We determined  $(2,90)$ -islands. The corresponding subnetwork has 470137 nodes, 307472 arcs and for different  $k$ :  $C_2 = 187610$ ,  $C_5 = 8859$ ,  $C_{30} = 101$ ,  $C_{50} = 30$ , ... islands.

## Rolex

[1]	0	139793	29670	9288	3966	1827	997	578	362	250
[11]	190	125	104	71	47	37	36	33	21	23
[21]	17	16	8	7	13	10	10	5	5	5
[31]	12	3	7	3	3	3	2	6	6	2
[41]	1	3	4	1	5	2	1	1	1	1
[51]	2	3	3	2	0	0	0	0	0	1
[61]	0	0	0	0	1	0	0	2	0	0
[71]	0	0	1	1	0	0	0	1	0	0
[81]	2	0	0	0	0	1	2	0	0	7



# Distribution of island size


Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores





# Main path and main island in US Patents

Nber, US Patents;  $n = 3774768$ ,  $m = 16522438$



Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores



V. Batagelj

Rnet, cohesion





# Main island Liquid crystal display

Rnet, cohesion

V. Batagelj

Islands

Cores

Generalized cores

Table 1: Patents on the liquid-crystal display

patent	date	author(s) and title
474969	Mar 14, 1994	Yvonne: Infrared light-polarizing sheet and the use and the formation of same
282562	Jan 29, 1984	Winder, et al. Rotation of aromatic carbonds
332485	Jun 20, 1987	Williams: Electro-optical elements utilizing an organic isotropic compound
363649	Jan 18, 1972	Josephson: Preparation of polymeric aromatic compounds
366498	May 30, 1972	Mokkanta, et al. Liquid crystal thermal imaging system having an undistorted image on a distorted background
375797	Jul 11, 1972	Rubin: Liquid crystal composition and device
380755	Sep 10, 1972	Orland: Check with digital display
380755	Oct 10, 1972	Wysocil: Electro-optic systems in which an electrophotoconductive or dipole material is dispersed throughout a liquid crystal to reduce the turn-off time
386498	May 8, 1973	Propoy: Display device utilizing liquid crystal light modulation
3757289	Oct 23, 1973	Aronson, et al. Class of stable trans-ethane compounds, some displaying nematic mesophase at or near room temperature and others in a range up to 100°C
373747	Nov 20, 1973	Stolzenberger: Substituted azo benzene compounds
379526	Mar 5, 1974	Bell, et al. Nematic liquid material which exhibits the Kerr effect at isotropic temperatures
379649	Mar 12, 1974	Bell, et al. Electro-optical liquid modulation and utilizing a nematic liquid material which exhibits the Kerr effect at isotropic temperatures
382140	Mar 18, 1975	Randerson, et al. Liquid crystalline compositions and method
3876284	Apr 8, 1975	Tomkinson, et al. Use of nematic liquid crystalline substances
3881806	May 6, 1975	Sanki: Electro-optical display device
3903387	Jun 24, 1975	Debnath, et al. Phase control of the voltage applied to opposite electrodes for a cholesteric to nematic phase transition display
3947275	Nov 30, 1976	Gray, et al. Liquid crystal materials and device
3954022	May 4, 1976	Stolzenberger: Liquid compositions having high dielectric anisotropy and display device incorporating same
3980712	Jun 1, 1976	Khandekar, et al. Liquid crystal compositions
4002586	Aug 17, 1976	Oh: Low voltage actuated field effect liquid crystal compositions and method of synthesis
4008824	Dec 28, 1976	Stolzenberger: Modified nematic compounds with positive dielectric anisotropy
4011173	Mar 8, 1977	Stolzenberger: Liquid crystal compositions and electro-optic device incorporating them
4013502	Mar 22, 1977	Stolzenberger: Liquid crystal compositions and electro-optic device incorporating them
4017446	Apr 12, 1977	Finkel, et al. Perylene-based 4-alkyl-4'-phenoxybenzophenone method for preparing same and liquid crystal compositions incorporating them
4022555	Jun 14, 1977	Rose, et al. Novel liquid crystal compounds and electro-optic device incorporating them
4025720	Jun 28, 1977	Bilson, et al. Electro-optic device
4037789	Mar 7, 1978	Goaj, et al. Optically active cyano-biphenyl compounds and liquid crystal materials containing them
4062428	Apr 4, 1978	Hsu: Liquid crystal composition and method

Table 2: Patents on the liquid-crystal display

patent	date	author(s) and title
408799	Apr 11, 1978	U.S. Nematic liquid crystal compositions
4113647	Sep 12, 1978	Coutts, et al. Liquid crystalline materials
4118232	Oct 3, 1978	Krause, et al. Liquid crystalline materials of reduced viscosity
4137052	Dec 19, 1978	Edman, et al. Liquid crystalline cyclohexane derivatives
4144163	Apr 17, 1979	Gray, et al. Optically active liquid crystalline and liquid crystal devices containing them
4154087	May 15, 1979	Edman, et al. Liquid crystalline benzophenonebiphenyl derivatives
4155294	Oct 23, 1979	Coutts, et al. Liquid crystal compounds
4196130	Apr 15, 1980	Bolke, et al. Liquid crystal materials
4202730	May 13, 1980	Sato, et al. Nematic liquid crystalline materials
4229215	Oct 21, 1980	Krause, et al. Liquid crystalline cyclohexane derivatives
4263052	Apr 14, 1981	Gray, et al. Liquid crystal compounds and materials and devices containing them
4299095	Sep 22, 1981	Kashe, Ester compound
4303434	Oct 6, 1981	Drostner, et al. Liquid crystal compounds
4320252	Nov 24, 1981	Edman, et al. Fluorophenylcyclohexanes, the preparation thereof and their use as components of liquid-crystal diisotrope
4330426	May 18, 1982	Edman, et al. Cyclohexylbiphenyls, their preparation and use in diisotropic and electro-optical display elements
4340498	Jul 20, 1982	Sagstrom: Halogenated ester derivatives
4349452	Nov 4, 1982	Omura, et al. Cyclohexylcyclohexanes
4357378	Nov 9, 1982	Carr, et al. Liquid crystal compounds containing an alkylic ring and exhibiting a low dielectric anisotropy and liquid crystal materials and device incorporating such compounds
4363494	Nov 30, 1982	Omura, et al. Anisotropic cyclohexyl cyclohexylbiphenyl ethers
4368132	Jan 13, 1983	Omura, Anisotropic compounds with negative or positive D <sub>11</sub> -anisotropy and low optical anisotropy
4387038	May 31, 1983	Krause, et al. Liquid crystalline naphthalene derivatives
4387038	Jun 7, 1983	Finkel, et al. 4-(trans-4'-alkylcyclohexyl) benzoic acid 4'-oxo-C <sub>6</sub> H <sub>4</sub> OBiphenyl esters
4387039	Jun 7, 1983	Sagstrom, et al. Trans-4-(trans-4'-alkylcyclohexyl)cyclohexyl esters and 4'-oxo-biphenyl esters
4400293	Aug 23, 1983	Honer, et al. Liquid crystalline cyclohexyl derivatives
4415470	Nov 15, 1983	Edman, et al. Liquid crystalline fluoro-containing cyclohexylbiphenyls and diisotropic and electro-optical display elements based thereon
4419063	Dec 6, 1983	Prud'homme, et al. Liquid crystalline cyclohexylbenzofuran derivatives
4422924	Dec 27, 1983	Sagstrom, et al. Liquid crystal benzene derivatives
4452443	Jun 19, 1984	Falkes, et al. Nematic halogen Compound
4456712	Jun 26, 1984	Sagstrom, et al. Cyclohexylcyclohexane compositions
4467279	Jul 17, 1984	Petrillo, et al. Liquid crystal materials
4472283	Sep 18, 1984	Sagstrom, et al. High temperature liquid crystal substances of low range and liquid crystal compositions containing the same
4472292	Sep 18, 1984	Falkes, et al. Nematic liquid crystalline compounds
4480117	Oct 30, 1984	Takata, et al. Nematic liquid crystalline compounds
4502074	Mar 5, 1985	Sagstrom, et al. High temperature liquid-crystalline ester compounds
4510809	Apr 9, 1985	Edman, et al. Cyclohexane derivatives

Table 3: Patents on the liquid-crystal display

patent	date	author(s) and title
4519414	Apr 30, 1985	Gray, et al. 4-(trans-4'-alkylcyclohexyl)-4'-oxo-C <sub>6</sub> H <sub>4</sub> OBiphenyl ester derivatives and liquid crystal mixtures
4526704	Jul 2, 1985	Petrillo, et al. Multisite liquid crystal esters
4529961	Nov 5, 1985	Petrillo, et al. Liquid crystalline esters and mixtures
4535121	Dec 10, 1985	Takata, et al. Nematic liquid crystalline compounds
4538262	Apr 22, 1986	Petrillo, et al. Phenoxyethanes
4531961	Nov 11, 1986	Petrillo, et al. Novel liquid crystal mixtures
4538396	Dec 23, 1986	Petrillo, et al. Benzoxiranes
4557089	Apr 14, 1987	Sato, et al. Substituted propanones
4565952	Apr 21, 1987	Furuta, et al. Ethane derivatives
4601531	Sep 22, 1987	Balkwill, et al. Disubstituted ethanes and their use in liquid crystal materials and devices
4704227	Nov 3, 1987	Krause, et al. Liquid crystal compounds
4706020	Nov 24, 1987	Petrillo, et al. Novel liquid crystal mixtures
4710315	Dec 1, 1987	Schub, et al. Anisotropic compounds and liquid crystal mixtures therefrom
4712197	Dec 15, 1987	Edman, et al. Nitrogen-containing heterocyclic compounds
4718022	Jan 12, 1988	Wahlberg, et al. Cyclohexane derivatives
4721267	Jan 26, 1988	Yoshizawa, et al. Liquid crystal device
4724144	Apr 21, 1988	Edman, et al. Nitrogen-containing heterocyclic compounds
4726505	Sep 13, 1988	Buchele, et al. Liquid crystalline compounds
4736579	Jan 3, 1989	Vankar, et al. 2,2'-difluoro-4-alkoxy-4'-hydroxydiphenyl ether and their derivatives, their production process and their use in liquid crystal display devices
4736579	Jan 10, 1989	Gray, et al. Cyclohexane derivatives and liquid crystal compositions containing same
4739233	Jan 11, 1989	Krause, et al. Nitrogen-containing heterocyclic esters
4823462	May 23, 1989	Clark, et al. Liquid crystal devices
4875747	Dec 11, 1989	Wolke, et al. Liquid crystal display element
4907349	Nov 1989	Chen, et al. Active matrix screen for the color display of television pictures, control system and process for producing said screen
5010898	May 21, 1991	Imura: Liquid crystal display device with a birefringent compensator
5018909	May 21, 1991	Ohira: Liquid crystal element with improved contrast and method
5122295	Nov 16, 1992	Wolke, et al. Matrix liquid crystal display
5124824	Jan 23, 1992	Komaki, et al. Liquid crystal display device comprising a retardation compensation layer having a maximum principal refractive index in the thickness direction
5171409	Dec 15, 1992	Hirata, et al. Liquid-crystal matrix display
5182673	Feb 1, 1994	Hirata, et al. Liquid-crystal display with grayed regions between terminal groups
5285338	Mar 3, 1994	Wolke, et al. Superwide liquid-crystal display
5374374	Dec 20, 1994	Wolke, et al. Superwide liquid-crystal display
5438877	Apr 6, 1996	Wolke, et al. Superwide liquid-crystal composition
5525316	Sep 10, 1996	Ichikawa, et al. Liquid crystal display having adjacent elements not set up in height
5604282	Nov 4, 1997	Schlagheckl, et al. Liquid crystal composition
5626814	Nov 4, 1999	Maher, et al. Liquid crystal compositions and liquid crystal display elements

V. Batagelj

Rnet, cohesion



# Word clouds for LCD island and foam island


Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores





# The Edinburgh Associative Thesaurus

$n = 23219$ ,  $m = 325624$ , transitivity weight





Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores



V. Batagelj

Rnet, cohesion





# Dense groups

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

Several notions were proposed in attempts to formally describe dense groups in graphs.

*Clique* of order  $k$  is a maximal complete subgraph (isomorphic to  $K_k$ ),  $k \geq 3$ .

$s$ -plexes, LS sets, lambda sets, cores, . . .

For all of them, except for cores, it turned out that they are difficult to determine.



# Cores and generalized cores


Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores



The notion of core was introduced by Seidman in 1983. Let  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$  be a graph. A subgraph  $\mathcal{H} = (W, \mathcal{E}|W)$  induced by the set  $W$  is a *k-core* or a *core of order k* iff  $\forall v \in W$  :  $\deg_{\mathcal{H}}(v) \geq k$ , and  $\mathcal{H}$  is a maximal subgraph with this property. The core of maximum order is also called the *main* core.

The *core number* of node  $v$  is the highest order of a core that contains this node. The degree  $\deg(v)$  can be: in-degree, out-degree, in-degree + out-degree, etc., determining different types of cores.



# Properties of cores

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

From the figure, representing 0, 1, 2 and 3 core, we can see the following properties of cores:

- The cores are nested:  $i < j \implies \mathcal{H}_j \subseteq \mathcal{H}_i$
- Cores are not necessarily connected subgraphs.

An efficient algorithm for determining the cores hierarchy is based on the following property:

*If from a given graph  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$  we recursively delete all nodes, and edges incident with them, of degree less than  $k$ , the remaining graph is the  $k$ -core.*



# 6-core of Krebs Internet industries


Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores





# Generalized cores

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

The notion of core can be generalized to networks. Let  $\mathcal{N} = (\mathcal{V}, \mathcal{E}, w)$  be a network, where  $\mathcal{G} = (\mathcal{V}, \mathcal{E})$  is a graph and  $w : \mathcal{E} \rightarrow \mathbb{R}$  is a function assigning values to edges. A *node property function* on  $\mathbf{N}$ , or a *p-function* for short, is a function  $p(v, U)$ ,  $v \in \mathcal{V}$ ,  $U \subseteq \mathcal{V}$  with real values. Let  $N_U(v) = N(v) \cap U$ . Besides degrees and (corrected) clustering coefficient, here are some examples of *p-functions*:

$$p_S(v, U) = \sum_{u \in N_U(v)} w(v, u), \text{ where } w : \mathcal{E} \rightarrow \mathbb{R}_0^+$$

$$p_M(v, U) = \max_{u \in N_U(v)} w(v, u), \text{ where } w : \mathcal{E} \rightarrow \mathbb{R}$$

$$p_t(v, U) = \frac{|\mathcal{L}(U) \cap \mathcal{L}(K(N^+(v)))|}{|\mathcal{L}(K(N^+(v)))|}$$

$$p_k(v, U) = \text{number of cycles of length } k \text{ through node } v \text{ in } (U, \mathcal{E}|U)$$

The subgraph  $\mathcal{H} = (C, \mathcal{E}|C)$  induced by the set  $C \subseteq \mathcal{V}$  is a *p-core at level*  $t \in \mathbb{R}$  iff  $\forall v \in C : t \leq p(v, C)$  and  $C$  is a maximal such set.



# Additional $p$ -functions

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

relative density

$$p_{\gamma}(v, \mathcal{C}) = \frac{\deg(v, \mathcal{C})}{\max_{u \in N(v)} \deg(u)}, \text{ if } \deg(v) > 0; 0, \text{ otherwise}$$

diversity

$$p_{\delta}(v, \mathcal{C}) = \max_{u \in N^+(v, \mathcal{C})} \deg(u) - \min_{u \in N^+(v, \mathcal{C})} \deg(u)$$

average weight

$$p_a(v, \mathcal{C}) = \frac{1}{|N(v, \mathcal{C})|} \sum_{u \in N(v, \mathcal{C})} w(v, u), \text{ if } N(v, \mathcal{C}) \neq \emptyset; 0, \text{ otherwise}$$



# Generalized cores algorithms

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

The function  $p$  is *monotone* iff it has the property

$$C_1 \subset C_2 \Rightarrow \forall v \in \mathcal{V} : (p(v, C_1) \leq p(v, C_2))$$

The degrees and the functions  $p_S$ ,  $p_M$  and  $p_k$  are monotone. For a monotone function the  $p$ -core at level  $t$  can be determined, as in the ordinary case, by successively deleting nodes with value of  $p$  lower than  $t$ ; and the cores on different levels are nested

$$t_1 < t_2 \Rightarrow \mathcal{H}_{t_2} \subseteq \mathcal{H}_{t_1}$$

The  $p$ -function is *local* iff  $p(v, U) = p(v, N_U(v))$ .

The degrees,  $p_S$  and  $p_M$  are local; but  $p_k$  is **not** local for  $k \geq 4$ .

For a local  $p$ -function an  $O(m \max(\Delta, \log n))$  algorithm for determining the  $p$ -core levels exists, assuming that  $p(v, N_C(v))$  can be computed in  $O(\deg_C(v))$ .

For details see the [paper](#).



# Cores and generalized cores / Pajek commands

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

```
File/Network/Read [Geom.net]
Network/Create Partition/k-Core/All
Info/Partition
Operations/Network+Partition/Extract Subnetwork [13-*]
Draw/Network+First Partition
Layout/Energy/Kamada-Kawai
Options/Values of lines/Similarities
Layout/Energy/Kamada-Kawai
Operations/Network+Partition/Extract Subnetwork [21]
Draw/Network
Layout/Energy/Kamada-Kawai
Options/Values of lines/Forget
Layout/Energy/Kamada-Kawai
[select Geom.net]
Network/Create Vector/Generalized Cores/Sum/All
Info/Vector
Vector/Make Partition/by Intervals/Selected Thresholds [4]
Info/Partition
Operations/Network+Partition/Extract Subnetwork [2]
Draw/Network
Options/Values of lines/Similarities
Layout/Energy/Fruchterman-Reingold
```



# Cores of orders 10–21 in Computational Geometry


Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores



V. Batagelj

Rnet, cohesion



# $p_S$ -core at level 46 of Geombib network

Rnet,  
cohesion

V. Batagelj

Islands

Cores

Generalized  
cores

